

# 2X BLEND DOSING

Tesamorelin 10mg / Ipamorelin 5mg = 15mg Total

## PROVIDER GUIDE

The chart below shows various dosing options coupled with the amount (in mg) and frequency for each dose. Dosing should be determined by the physician on an individual patient basis.

### Reconstitution Instructions

For our Tesamorelin (10mg)/Ipamorelin (5mg), add 1mL of bacteriostatic water → **0.15mg/unit** concentration

Dose	Volume Drawn	Tesamorelin	Ipamorelin
1.5mg	10 units	1mg	0.5mg
2.7mg	18 units	1.8mg	0.9mg
3.6mg	24 units	2.4mg	1.2mg
4.5mg	30 units	3mg	1.5mg



#### Suggested Dose

1.5mg–4.5mg  
(10 to 30 units)



#### Administration

**Method**  
Subcutaneous



#### Dose Frequency

5 days on, 2 days off



#### Blend Benefits

Best for anti-aging mixed  
with heavy weight loss



#### Primary Benefit

Fat loss and lean muscle



#### Notes

Fast 45 minutes before  
and after injection

# 2X BLEND DOSING

Tesamorelin 5mg / Ipamorelin 5mg = 10mg Total

## PROVIDER GUIDE

The chart below shows various dosing options coupled with the amount (in mg) and frequency for each dose. Dosing should be determined by the physician on an individual patient basis.

### Reconstitution Instructions

For our Tesamorelin (5mg)/Ipamorelin (5mg), add 1mL of bacteriostatic water → **0.1mg/unit** concentration

Dose	Volume Drawn	Tesamorelin	Ipamorelin
1.5mg	15 units	0.75mg	0.75mg
2.5mg	25 units	1.25mg	1.25mg
3.5mg	35 units	1.75mg	1.75mg
4.5mg	45 units	2.25mg	2.25mg



#### Suggested Dose

1.5mg–4.5mg  
(15 to 45 units)



#### Administration

##### Method

Subcutaneous



#### Dose Frequency

5 days on, 2 days off



#### Blend Benefits

Best for anti-aging mixed  
with heavy weight loss



#### Primary Benefit

Fat loss and lean muscle



#### Notes

Fast 45 minutes before  
and after injection