



**Lyophilized Tirzepatide**  
**Dosing & Reconstitution**  
**Guidelines**  
 For Providers

Below are charts showing various dosing options coupled with the amount of Bacteriostatic Water added. Dosing should be determined by the physician on an individual patient basis.

**Provider Guide: Dosing for Injectable Lyophilized Tirzepatide**

Amount In Vial	Bac Water In	Amount Per Unit	Recommended Dose
10 mg	1 mL	0.1 mg per unit	Titrate up from 2.5mg to 10mg (Titrate up from 25 to 100 units)
30 mg	3 mL	0.1 mg per unit	Titrate up from 2.5mg to 15mg (Titrate up from 25 to 150 units)
50 mg	3 mL	0.17 mg per unit	Titrate up from 2.5mg to 15mg (Titrate up from 15 to 90 units)
60 mg	3 mL	0.2 mg per unit	Titrate up from 2.5mg to 15mg (Titrate up from 12.5 to 75 units)

**More Information**

Frequency	Notes
1x per week	Injection type: Subcutaneous Same day every week / starting dose is 2.5mg



**Bacteriostatic Water**

This solution allows for the safe storage of reconstituted peptides, like Tirzepatide GLP-2, in refrigeration for up to 60 days, ensuring that the peptide remains effective and uncontaminated while maintaining peptide integrity and potency.